

by Kenitiro Suguio^{1,2}, Alethéa Ernandes Martins Sallun², and Emílio A. A. Soares^{2,3}

Quaternary: “Quo Vadis”?

1 Centro de Pós-Graduação, Pesquisa e Extensão (CEPPE), Universidade Guarulhos (UnG), Praça Tereza Cristina, 01 – Guarulhos – SP – Brazil – 07023-070. e-mail: kenitirosuguio@hotmail.com

2 Programa de Pós-Graduação em Geologia Sedimentar - Instituto de Geociências, Universidade de São Paulo, Rua do Lago, 562 – São Paulo – SP – Brazil – 05508-080. e-mail: aletheamartins@hotmail.com

3 Departamento de Geociências, Universidade Federal do Amazonas, Av. Gal. Rodrigo O.J. Ramos, 3000, Manaus – AM – Brazil - 69077-000 e-mail: easoares@usp.br

The Quaternary, like the long-abandoned terms Primary, Secondary and Tertiary, is a very ambiguous word, whose chronological meaning is not very well defined. Its usage spread very quickly, perhaps due to its more-or-less close relationship with the human history and with the latest glaciations. However, the oldest of human fossil record, according to recent discoveries in Africa (Djourab Desert in Tchad), is about 7 million years old, rather than 2 million years as suggested by fossils at Olduvai. Moreover, for some time now, it has been known that the first Quaternary-type glaciations occurred in several regions of the Earth between 2.5 and 3 million years ago, that is prior to the Pliocene-Pleistocene limit at 1.81 million years ago, as determined by the GSSP. These facts have led some researchers, mostly from the International Union of Geological Sciences (IUGS), and from the International Association for Quaternary Research (INQUA), to suggest changes in the subdivision of the Neocenozoic. Certainly, discussions on this subject will be raised at many scientific meetings in the near future until almost unanimous agreement can be attained. Some additional suggestions are presented here, as a contribution to resolution of these nomenclatorial issues.

Introduction

According to Phillips (1840), the present *Cenozoic Era* was subdivided in schemes still subjected to much discussion, until at least the middle of the 19th century, based on field relationships and/or on biological evolution. In 1760 Arduino, professor of the Padova University (Italy), used the word *Primary* for the oldest rocks followed by rocks of *Secondary*, during lithostratigraphical classification surveys of northern Italy mountainous region. The lower hills composed of gravelly, sandy and clayey sediments were attributed to the *Tertiary*. On the other hand, the term *Quaternary*, Desnoyers (1829) was proposed for alluvial and marine sediments containing the remains of associations of extant animals and plants, and overlying *Tertiary* deposits of the Paris Basin.

The use of the word *Quaternary*, even without more precise chronological definition, spread very rapidly particularly in the mapping of less consolidated superficial deposits. Ages attributed to *Quaternary* units on older geologic maps are very uncertain.

Based on relative similarity of fossil assemblages with the modern fauna, Lyell (1833) subdivided the *Tertiary* Period into the *Eocene*, *Miocene* and *Pliocene Epochs*. Apparently ignoring the previous proposal from Desnoyers (1829), Lyell named the post-Tertiary time as *Recent* (Fairbridge, 1968). The *Recent* Epoch was later renamed *Holocene Epoch* by Gervais (1867). Lyell never used the word *Quaternary*, but assumed that Desnoyer's concept was approximately equivalent to the time interval from the *Tertiary* Period *Newer Pliocene* through the *Recent*.

The *Neogene* and *Paleogene* Stages were introduced by Hörnes (1853) to subdivide Cenozoic deposits and were adopted by the International Commission on Stratigraphy (ICS). The *Paleogene* of Hörnes included the *Paleocene*, *Eocene* and *Oligocene Epochs* and the *Neogene*, the *Miocene*, *Pliocene*, and *Pleistocene Epochs*.

When the International Commission on Stratigraphy (ICS) and the International Association for Quaternary Studies (INQUA) decided to standardize the *Pliocene-Pleistocene* limit during the 1950s, there were three proposals for formal definition:

- Lyell's (1833) *Newer Pliocene* at approximately 1 My;
- Top of Olduvai palaeomagnetic subzone at about 2 My, and
- Near the Gauss-Matuyama palaeomagnetic polarity reversal at about 2.5 My.

Option (b) was chosen during the INQUA International Congress in 1982 (Moscow) by the INQUA Commission on Stratigraphy, acting as a subcommission of the International Commission on Stratigraphy (ICS); it was formally approved by this commission in 1983. According to this decision, the base of the Pleistocene (boundary stratotype), composed of deepwater marine deposits, would be located in Vrica, Sicily (Italy), near the top of Olduvai normal polarity palaeomagnetic subzone at about 1.8 My. It characterizes the time of greatest spreading of the mollusk *Arctica islandica* which was, in general, thought to be restricted to boreal waters of the interglacial stage. According to Aguirre & Pasini (1985), this boundary stratotype does not take into account the situation of the *Quaternary* in the chronostratigraphic scale. The stratotype was presented by Pasini & Colalongo (1997) and, moreover, was characterized in detail by stratigraphers from many countries from sedimentological, palaeoecological, biostratigraphical, biochronological and magnetostratigraphical viewpoints.

The word *Quaternary*, even without a formal definition, became very entrenched and amongst other peculiarities, traditionally has been correlated with glacial episodes of the Northern Hemisphere (Figure 1).

However, chronological evidence, based on marine isotopic records and on ice-drifted sediments of the Northern Atlantic Ocean, indicates that the most important increase in the volume of continental glaciers must have begun about 2.6 Ma. Because of this, this subject was reevaluated in 1998 by the Neogene Stratigraphy Commission of ICS and by the Quaternary Stratigraphic Commission of ICS-INQUA, which again resulted in the formal rejection of placement of the Pliocene-Pleistocene limit at the base of the Gelasian Stage. Its boundary-stratotype was maintained at base *Pleistocene* in Vrica, Sicily (Italy), dated at 1.806 Ma. Thus, if the *Quaternary* is defined on the basis of the most important oscillations in Northern Hemi-

PERIOD	EPOCH	MAGNETO-CLASSIFICATION	AGES X 10 ³ YEARS	¹⁸ O ISOTOPE STAGES	ALPINE GLACIATION (STAGE)	LAURENTIAN GLACIATION (STAGE)	ENGLAND GLACIATION (STAGE)	SCANDINAVIAN GLACIATION (STAGE)	NORTH EUROPEAN - PALYNOLOGICAL ZONES	ITALIAN GLACIATION							
QUATERNARY	Holocene	[Black bar]	1	1	Flandrian	Recent		Flandrian	Flandrian	Recent Sub-Atlantic Sub-Boreal Atlantic	Versilian						
			Late Pleistocene	[Black bar]	10	2	Later	Wisconsin Glacial	Devensian	Later	Weichselian	Transition Boreal Younger Dryas Allerød Older Dryas Bölling Oldest Dryas	Tyrrenian				
					20	3	Middle			Middle		Denekamp Hengelo					
					50	4	Early			Early		Odderade					
					5a	Ig. Riss-Würm						Sangamon Ig.		Ipswichian	Eemian		
	5c																
	100	5e															
	Middle Pleistocene	[Black bar]			[Black bar]	200	7			Riss Glacial		Illinoian Glacial		Wolstonian	Saalian		Crotonian
						300	9			Ig. Mindel-Riss		Yarmouthian Ig.		Hoxnian	Holsteinian		
						400	11			Mindel Glacial		Kansanian Glacial		Anglian	Elsterian		
						500	13			Günz-Mindel interglacial		Aftonian Interglacial		Cromerian*	Noordber.		
			600	15		Rosmalen.	Westerhov.										
			700	17													
			800	19		Waardenb.											
			900			Leedam											
1000				Bavel													
Early Pleistocene			[Black bar]	[Black bar]		1500		Günz Glacial	Nebraskan Gi.	Bavention	Menapian		Sicilian				
					Donau-Günz Ig.		Antian	Waalian		Emilian							
					Donau Glacial		Thurnian	Eburonian		Santerian							

Figure 1 Tentative correlation between the Quaternary Period and Northern Hemisphere glacial and interglacial (Ig) episodes (Geosciences Research Group of Japan, 1996).

sphere glacier volume, it began 800 thousand years before the Pleistocene Epoch base (Figure 2).

Some proposals for formalizing the Quaternary

Subdivision proposed by the International Commission on Stratigraphy (ICS)

The Cenozoic Era, with a total duration of about 65 My, would be subdivided into *Paleogene* (42 My) and *Neogene* (23 My) *Periods*, consisting of comparable time intervals (Figure 3). The *Tertiary* (about 63 My or more than 95% of the Cenozoic) and the *Quaternary* (about 2 My or less than 5% of the Cenozoic) *Periods* would be formally abandoned, but maintained informally, like the equally important unit term Precambrian.

The decision to abandon the word *Tertiary* follows in the same tradition that led to the suppression of *Primary* and *Secondary*: they are very ambiguous words. In the subdivision presented by Gradstein et al. (2004), *Quaternary* was also omitted, but *Pleistocene* and

Holocene were maintained with the *Pliocene-Pleistocene* boundary being set at 1.8 My and the *Pleistocene-Holocene* limit at 0.0115 My.

Subdivision of the International Association for Quaternary Research (INQUA)

Recently, Pillans (2004) emphasized the need for maintaining the *Quaternary*, as an international subdivision of the *Neogene Period* (Figure 4). According to this author, the *Quaternary* represents a word too important to be simply omitted from the Geologic Time Scale, as happened with the *Primary*, *Secondary* and, more recently, *Tertiary*. As one his justifications, the author states that the *Quaternary* represents a link between humankind and geology. Moreover, it would provide the needed umbrella-type protection for other disciplines correlatable with geosciences, as for example, archaeology, palaeopedology, palaeoclimatology, etc. Pillans proposed redefinition of the *Quaternary* as a Subperiod (or Subsystem) of the Neogene Period (or System), beginning at about 2.6 My and including at its base the Gelasian stage (Rio et al., 1998). The most important arguments in favor of this proposal are:

a) There is very strong support by INQUA members, who answered positively to the proposal for maintenance of the *Quaternary* as a formal stratigraphic unit.

b) Precedence already exists for introducing the Subperiod (or Subsystem) in the Geologic Time Scale as, for example, the Mississippian and Pennsylvanian Subperiods (or Subsystems) of the Carboniferous Period.

c) Disassociation of the base of the *Quaternary* from the Pliocene-Pleistocene boundary (1.8 My) would finish discussions about the position of this limit.

d) The majority of INQUA members seems to be favorable to the "longer" Quaternary (2.6 My), instead of the "shorter" Quaternary (1.8 My). This choice reflects the

CENOZOIC							
AGE (My)	Period	Epoch	Stage	AGE (My)			
0	Neogene	Pleistocene	Holocene	Late	1.8		
			Pliocene	L		Middle	
						Early	
		Pliocene	E	E		Gelasian	2.6
						Piacenzian	
						Zanclean	
		Miocene	E	E		Messinian	5.3
							7.3

Figure 2 Current span of the Quaternary without formal definition based on the most important oscillations in Northern Hemisphere glacier volumes (Ogg, 2004).

Epoch	Eon	System	Series	Stage	Age
Phanerozoic					
Cenozoic					
Neogene					
Pliocene					
Pleistocene					
Holocene					
Paleogene					
Oligocene					
Eocene					
Paleocene					
			Holocene		0.0115
			Pleistocene	Upper	0.126
				Middle	0.781
				Lower	1.806
			Pliocene	Gelasian	2.588
				Piacenzian	3.600
				Zanclean	5.332
			Miocene	Messinian	7.246
				Tortonian	11.608
				Serravallian	13.65
				Langhian	15.97
				Burdigalian	20.43
				Aquitanian	23.03
				Chattian	28.4±0.1
			Rupelian	33.9±0.1	
			Eocene	Priabonian	37.2±0.1
				Bartonian	40.4±0.2
				Lutetian	48.6±0.2
				Ypresian	55.8±0.2
			Paleocene	Thanetian	58.7±0.2
				Selandian	61.7±0.2
				Danian	65.5±0.3

Figure 3 Proposal of the International Commission on Stratigraphy (ICS) of the International Union of Geological Sciences (IUGS) published by Gradstein et al. (2004) and Lourens et al. (2004).

understanding of significance of continuity of the properties through time. For example, loess deposition in China became more intense and more extensive about 2.6 My ago, with quite different properties from the underlying “red clays” (Ding et al. 1997).

e) At about 2.6 My, deep sea oxygen isotope records show a series of cycles of growing glacial intensities, which are also associated with first record of more abundant North Atlantic Ocean glacial detrital sediments. For many researchers this would represent the advent of Quaternary glacial ages. This boundary corresponds to the transition from equinox precession to ecliptical obliquity as the dominant process in climatic forcings (Milankovitch, 1920).

According to the INQUA Stratigraphy and Chronology Commission, consultations about this subject must continue, during the International Union of Geological Sciences (IUGS) congresses, like that recently finished in Florence (Italy), as well as during the next INQUA International Congress in Cairns (Australia), to be held in 2007.

Present proposal

It is suggested here that, for the same reasons alleged for the abandonment of the terms Primary, Secondary and Tertiary, the word Quaternary could be eliminated from the Geological Time Scale as well, because of its extreme ambiguity. The word Holocene, which is commonly considered as synonymous of Recent or Post-glacial, could be also omitted (Figure 5).

In support of our suggestion, the following questions could be raised:

a) Why Post-glacial, if until now there is no irrefutable evidence that Pleistocene glaciations have really finished?




Figure 4 Present situation of the Quaternary and the proposal by Pillans (2004), changing Quaternary to Subperiod (or Subsystem) of the Neogene Period (or System).

b) Why couldn't the well-known mild climate of the Holocene (or Post-glacial) simply represent an interglacial stade, as many palaeoclimatologists think?

c) If the previous hypothesis is believable, isn't an imminent new glacial stade quite possible within the near future, in a few hundred or few thousand years?

In this case, the Pleistocene Epoch could be extended till the present. The Holocene, as a possible Pleistocene interglacial stade, does not deserve any formal title as an Epoch of Neogene Period (Suguio & Soares, 2004). On the other hand, the above mentioned uncertainties do not annul the possibility to consider the Quaternary as an informal chronostratigraphic unit, whose beginning could be located at the base of the Gelasian Stage, coincident with the commencement of the Pleistocene Epoch at about 2.6 My, thereby reducing the duration of Pliocene Epoch by 800 thousand years. This informal use of Quaternary would be somewhat analogous to that of Precambrian, at the other end of the time scale, maintained as such by the strength of tradition. However, we are aware that it remains to be seen if re-opening (for the third time in the last two decades) of

Epoch	Eon	System	Series	Stage	Age
Phanerozoic					
Cenozoic					
Neogene					
Pliocene					
Pleistocene					
Holocene					
Paleogene					
Oligocene					
Eocene					
Paleocene					
			Pleistocene	Upper	0.126
				Middle	0.781
				Lower	1.806
			Pliocene	Gelasian	2.588
				Piacenzian	3.600
				Zanclean	5.332
			Miocene	Messinian	7.246
				Tortonian	11.608
				Serravallian	13.65
				Langhian	15.97
				Burdigalian	20.43
				Aquitanian	23.03
				Chattian	28.4±0.1
			Rupelian	33.9±0.1	
			Eocene	Priabonian	37.2±0.1
				Bartonian	40.4±0.2
				Lutetian	48.6±0.2
				Ypresian	55.8±0.2
			Paleocene	Thanetian	58.7±0.2
				Selandian	61.7±0.2
				Danian	65.5±0.3

Figure 5 New proposal for the Neogene Period subdivision, composed by Miocene, Pliocene and Pleistocene, with omission of the Holocene Epoch.

the formal debate on re-positioning of the base Pleistocene would be fruitful.

Final considerations

The use of anthropological criteria in characterizing the last chapter (Pleistocene Epoch or "Quaternary") of historical geology makes the Geological Time Scale too anthropocentric, and apparently inconsistent, since mankind may well have appeared in the Miocene Epoch. On the other hand, humankind became conscious of its own existence only about 10 thousand years ago (at the beginning of the "Holocene"). At that time, mankind began to abandon instinctive wandering behaviour, like wild animals, and adopted more sedentary life. This change in lifestyle became possible with "domestication" of animals and plants for food.

The abandonment of Holocene and maintenance of "Quaternary", as an informal chronostratigraphic unit are proposed in this paper. The Quaternary would become synonymous with the Pleistocene Epoch, which began about 2.6 My. This proposal would safeguard the continuity of such important and traditional research organizations as INQUA and similar national associations, like the Brazilian Association of Quaternary Studies (ABEQUA), as well as of their on going multi and interdisciplinary studies. Finally, the authors believe that this proposal is the best one, because it is in perfect agreement with the Geological Timetable based on the Earth history and evolution.

The authors are grateful to Dr. Thomas Rich Fairchild for careful revision of the English text.

References

- Aguirre, E. and Pasini, G., 1985, The Pliocene-Pleistocene boundary: Episodes, v. 8, pp. 116 – 120.
- Arduino, G., 1760, A letter to Sig. Cav. Antonio Valisnieri: Nuova raccolta di opuscoli scientifici e filologici del padre Angiolo Calogierà (Venice), v. 6, pp. 142 – 143.
- Desnoyers, J., 1829, Observations sur un ensemble de dépôts marins plus récents que les terrains tertiaires du bassin de la Seine, et constituant une formation géologique distincte: précédées d'une aperçu de la non-simultanéité des bassins tertiaires: Annales Sciences Naturelles, Paris, v. 16, pp. 117 – 214, pp. 402 – 491.
- Ding, Z., Rutter, N.W. and Liu, T., 1997, The onset of extensive loess deposition around the G/M boundary in China and its palaeoclimatic implication: Quaternary International, v. 40, pp. 53 – 60.
- Fairbridge, R. (ed.), 1968, The Encyclopedia of Geomorphology: Reinhold, pp. 1295.
- Geosciences Research Group of Japan, 1996, Cyclopedia of Geosciences: Heibonsha, 2 v. (in Japanese), pp. 1443.
- Gervais, P., 1867, Zoologie et paléontologie générales: Nouvelles recherches sur les animaux vertébrés et fossiles, p. 262.
- Gradstein, F.M., Ogg, J.G., Smith, A.G., Bleeker, W. and Lourens, L.J., 2004, A new Geologic Time Scale, with special reference to Precambrian and Neogene: Episodes, v. 27, pp. 83 – 100.
- Hörnes, M., 1853, Mitteilung an Prof. Bronn Gerichtet: Wien, 3. Okt., 1853: Neues Jahrb. Mineral. Geol. Geogn. Petrefaktenkd., pp. 806 – 810.
- Lourens, L., Hilgen, F., Shackleton, N.J., Laskar, J. and Wilson, D., 2004, The neogene Period. *in* Gradstein, et al. eds, A Geologic Time Scale 2004. p. 409-440. Cambridge University Press, UK.
- Lyell, C., 1833, Principles of Geology: being inquiry how far the former changes of the earth's surface are referable to causes now in operation: London, John Murray, v. 3, pp. 398.
- Lyell, C., 1857, Supplement to the fifth edition: London, John Murray, pp. 40.
- Milankovitch, M., 1920, Théorie mathématique des phénomènes thermiques produits par la radiation solaire: Academie Yougoslave des Sciences et Arts, pp. 339.
- Ogg, J., 2004, Introduction to concepts and proposed standardization of the term "Quaternary": Episodes, v. 27, pp. 125– 126.
- Pasini, G. and Colalongo, M. L., 1997, The Pliocene- Pleistocene boundary-stratotype at Vrica, Italy, in J.A. Van Couvering, The Pleistocene boundary and the beginning of the Quaternary: Cambridge University Press, World and Regional Geology 9, pp. 15 - 45.
- Phillips, B., 1840, Paleozoic Series, in Long, G., ed, The Penny Cyclopaedia of the Society for the Diffusion of Useful Knowledge: London, Charles Knight, v.17, pp. 153 – 154.
- Pillans, B., 2004, Proposal to redefine the Quaternary, in Revision of the Geological Time Scale: Quaternary Perspectives, v.14, pp. 125.
- Rio, D., Sprovieri, R., Castradori, D. and Di Stephano, E., 1998, The Gelasian Stage (Upper Pliocene): A new unit of the global standard chronostratigraphic scale: Episodes, v. 21, pp. 82 – 87.
- Suguio, K., and Soares, E.A.A., 2004, Período Quaternário: "Quo vadis"?: XLII Congresso Brasileiro de Geologia, Araxá, Brazil, Anais, C753.

Kenitiro Suguio is professor emeritus of geology at the University of São Paulo, Brazil, where he remains as a lecturer and advisor in the graduate course on Sedimentary Geology. During the last 25 years he has been active participant of IGCP and IGBP projects on Quaternary Geology. He received the Brazilian National Book Prize in 1993 and the commendation of the National Order of Scientific Merit in 1998. He is also full professor at CEPPE (Centro de Pós-graduação, Pesquisa e Extensão), of Guarulhos University (State of São Paulo).



Alethéa Ernandes Martins Sallun is a graduate student in the Sedimentary Geology program of the Geosciences Institute, University of São Paulo. Her PhD research is on the Cenozoic alluvial-colluvial deposits in southeastern Brazil, supported financially by FAPESP.



Emílio Alberto Amaral Soares is professor of sedimentology in the Geoscience Department, Amazon Federal University, Brazil, and currently in the doctorate program at the Geoscience Institute, University of São Paulo. His dissertation subject is the evolution of Cenozoic deposits of the Amazon Basin, particularly as related to the latest depositional and climatic events of the Quaternary and their relation with Andean tectonics.

